

REFRACTION IN CONVEX AND **CONCAVE SPHERES**

Fig. 1: Refraction in a convex lens



A convexlens (pluslens)isliketwoprisms placed base to base. Light passing through a convex lens is converged. Convex lenses are used to treat presbyopia, hypermetropia and aphakia. If parallel light is brought to a focus at 1 metre the lens is said tohave 1 dioptreofpower. If the focus is at 1/2 metre, 2 dioptres, and at 1/3 metre, 3 dioptres(Fig.1).

Fig. 2: Refraction in a concave lens



A concave lens (minus lens) is like two prisms placed apex to apex. Light passing through a concavelensis diverged. Concave lenses are used to treat myopia (Fig. 2).

REFRACTION OF THE EYE

Fig. 3: Refraction of the eye



Light entering the eye is converged at the cornea and then the lens acts as a focusing mechanism to converge the light to a point on the retina, so that the object is seen clearly (Fig. 3).

Fig. 4: Refraction of the eye when the object is close: accommodation




If the object is close to the eye then the lens changes shape so that the light rays can still be focused on the retina. This is called accommodation. (The ability to accommodate begins to fail after the age of 40, so that spectacles are need for close work). (Fig. 4).

Fig. 5: Refraction in the eye: myopia



In myopia light rays are focused in front of the retina, so that a minus concave lens is needed to diverge the rays (Fig.5).

Fig.6: Refraction in the eye: hypermetropia (hyperopia)



In hypermetropia, it is the opposite, the rays are focused behind the retina so that a plus convexlensisneeded(Fig.6).

REFRACTION IN CYLINDERS

A cylindrical lens has different powers in the vertical and horizontal axes. Thus, light passing through a cylindrical lens does not focus at one point, but forms two foci, one for the horizontal and the other for the vertical. Cylindrical lenses may be convex, concave or mixed. They are used to treat astigmatism.

Illustrations and Text Reproduced by **Kind Permission**

From: Hanyane: A Village Struggles for Eye Health Erika Sutter, Allen Foster, Victoria Francis

Published by the International Centre for Eye Health. (Also available in French)

Available from:

International Resource Centre, ICEH 11-43 Bath St., London, EC1V 9EL

Price: £5/\$9

Postage and Packing: Overseas: add £3/\$5 (surface) or £5/\$9 (airmail) for each item UK: add £2 for each item

Payment is by bankers' order or cheque (drawn on US\$ or UK£ bank only) made out to University College London.

☆ ☆ ☆

A Manual of Refraction for Eye Health Workers

Developed by: Mr. Steve Mayer Optometrist, CBM International

This excellent Manual of Refraction has 45 pagesandincludesthefollowingsections:

- basic optics
- pinhole
- accommodation
- neutralisation
- transposition
- best sphere
- crossed cylinder
- cycloplegia
- optical lenses
- visual acuity
- refractive errors
- presbyopia focimetry
- spherical
- equivalent
- retinoscopy
- refraction
- basic spectacle dispensing.

Price: £5/US\$9 + post and packing

Post and Packing: Please add £5/\$9 (surface) or £9/\$16 (air)

Payment is by bankers' order or cheque (drawn on US\$ or UK£ bank only) made out to University College London.

Pleasesendyourorderwithpaymentto: **International Resource Centre, ICEH** Institute of Ophthalmology, 11-43BathSt.,LondonEC1V9EL,UK

Community Eye Health Courses 2000/2001

- MSc in Community Eye Health 1 year (Sept. 2000 Sept. 2001)
- Diploma in Community Eye Health 6 months (Sept. 2000 Mar. 2001)
- Certificate in Community Eye Health 3 months (Sept. Dec. 2000)
- Certificate in Planning for Eye Care 3 months (Jan. Mar. 2001)
- Short courses 1–4 weeks (on-going)
- Intensiveoneweekcourse: Planning EyeProgrammes (26–30June 2000)

Enquiries: Courses Promotions Officer, International Centre for Eye Health, 11-43 Bath Street, LONDON, EC1V 9EL, United Kingdom. Fax: +44 (0) 207 608 6950; E-mail:j.hoskin@ucl.ac.uk